



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## MECHANICS.

205. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

Given two points  $A$  and  $B$  not in the same horizontal nor in the same vertical line; to find the path from  $A$  to  $B$  along which a particle will slide from rest under the force of gravity alone so that the average velocity along the curve shall be a maximum.

Solution by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

Let the particle start from rest at the point  $(x_1, y_1)$ , where the axis of  $x$  is vertical and the axis of  $y$  horizontal. Then

$$v = \frac{ds}{dt} = \sqrt{2g(x-x_1)} = \frac{dx}{dt} \sqrt{1 + \left(\frac{dy}{dx}\right)^2}$$

Since  $v$  is a maximum,  $t$  is a minimum.

$$\therefore t = U = \int_{x_1}^{x_2} \frac{\sqrt{1 + (dy/dx)^2}}{\sqrt{2g(x-x_1)}} dx = \text{minimum.}$$

By Calculus of Variations, since the expression in the right hand member does not contain  $y$  explicitly, the differential of this expression is equal to a constant.

$$\therefore \frac{dy/dx}{\sqrt{2g(x-x_1)}} \frac{1}{\sqrt{1 + (dy/dx)^2}} = C.$$

$$\therefore \left(\frac{dy}{dx}\right)^2 = C^2 \left[1 + \left(\frac{dy}{dx}\right)^2\right] [2g(x-x_1)].$$

$$\therefore \frac{dy}{dx} = \pm \frac{C\sqrt{2g(x-x_1)}}{\sqrt{1 - 2gC^2(x-x_1)}},$$

the differential equation of a cycloid. This is Bernoulli's famous problem and is solved in almost every work on the Calculus of Variations as well as Dynamics.

206. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

A rigid square  $ABDC$  made by smooth wires is fixed with  $A$  vertically above  $D$ . Two small equal spherical elastic beads slide down  $BD$ ,  $CD$ , starting simultaneously from  $B$  and  $C$ . Find the ratio of their velocities of approach and separation at  $D$ , and how far they will separate after impact.